

Top Gun Kennel Vaccination Program

We do not dictate or recommend vaccination programs to our clients. We would strongly encourage that you establish a Vet-Client relationship with a local Vet practice in your area who can tailor your program to what is best in your geographical area. We have put our current immunization program in writing to help our customers and their veterinarians understand what we have done prior to the purchase of the puppy or adult dogs. This program has been used on our adult dogs and is required with a signed health certificate stating the immunization status of the dogs that we train here. Accurate immunization records must be filed on each animal for future use. Most states are now requiring this for interstate travel.

Week 1. Dock tails and remove dewclaws. Treat open wounds with antibiotics

Week 2. Deworm puppies with paste or liquid dewormer like Strongid or Nemec.

Week 4. Start the weaning process. Feed moistened puppy food and fresh water often.

Week 5. Discontinue placing the Bitch with the litter. Monitor stools, food consumption, and individual performance on litter. Place dry puppy food as free choice but continue feeding moistened puppy food and fresh water. Deworm with Drontal dewormer.

Week 6. Immunize Canine with Da2PP+CVK (Distemper, Adenovirus Type 2, Coronavirus, Parinfluenza, and Parvovirus). Modified live and killed virus, subcutaneous. Deworm with Fenbendazole dewormer.

Weeks 9 & 12. Immunize Canine with Da2LPP +CVK (Distemper, Adenovirus Type 2, Coronavirus, Parinfluenza, Parvovirus, Leptospira Bacterin). Immunize Canine Bordetella Bronchiseptica intranasal. Deworm with Fenbendazole dewormer.

Week 16. Immunize Canine Distemper, Adenovirus Type 2, Coronavirus, Parinfluenza, and Parvovirus, and Leptospira Bacterin Modified Live Virus subcutaneous and Immunize Canine Bordetella Bronchiseptica subcutaneous. Deworm with Fenbendazole dewormer. Immunize 1-year Rabies vaccination.

Adult Dogs. Annual Da2LPP+CVK and rotate intranasal and subcutaneous Canine Bordetella Bronchiseptica. Monthly prevention of Heartworm and external parasites, monitor sample stools quarterly for internal parasites and treat accordingly.

Vaccination for Borrelia Burgdorferi is recommended in areas of high tick infestation. Consult your veterinarian regarding this particular vaccination program!

Heartworm preventative should be administered by April 1 of each year through November or after the first killing frost occurs.

“The main reason to immunize your dog properly is because others don’t”

Top Gun Kennel Veterinarian:
Ross Vet Clinic – Dr Kathy Ross
3251 North Marion Road
Marion, IA 52302
(319) 373-1131